

Topology and its Applications 52 (1993) 109–120
North-Holland

109

Representing homology classes of almost definite 4-manifolds

Gao Hongzhu

Institute of Systems Science, Academia Sinica, Beijing 100080, People's Republic of China

Received 26 November 1990

Revised 10 April 1992

Abstract

Gao Hongzhu, Representing homology classes of almost definite 4-manifolds, *Topology and its Applications* 52 (1993) 109–120.

For simply connected 4-manifolds $\mathbb{C}P^2 \# m\overline{\mathbb{C}P}^2$, the representing problem of homology classes by embedded spheres is studied. A nonrepresentability condition is given. By using properties about automorphisms of unimodular quadratic forms over integers, some representability results are obtained.

Keywords: 4-manifold; Embedded sphere.

AMS (MOS) Subj. Class.: Primary 57N13.

1. Introduction

The content of this paper is organized as follows: first, using the result of Wall [5], we analyse a basic property about automorphisms of certain unimodular quadratic forms over integers. This is given mainly for later argument. Secondly, we investigate the problem about representing homology classes in almost definite 4-manifolds $\mathbb{C}P^2 \# m\overline{\mathbb{C}P}^2$ and obtain some nonrepresentability results about homology classes with positive square (Theorem 3.1). Combining this with the first part, we can determine such classes completely for $m = 2$ and 3 (Theorems 3.3 and 3.4). As a consequence, we may deduce the theorems of Kuga [2] and Lawson [3] from our results. Finally, some positive results are given (Theorem 3.7).

Correspondence to: Dr. Gao Hongzhu, Department of Mathematics, Beijing Normal University, Beijing 100875, People's Republic of China.

2. Action of orthogonal groups

Let X be a free Abelian group with a symmetric bilinear map ϕ of $X \times X$ to the integers, with determinant ± 1 , write $x \cdot y$ for $\phi(x, y)$. An automorphism of the bilinear map is an automorphism of group X , such that $\phi(Tx, Ty) = \phi(x, y)$ for all $x, y \in X$. These automorphisms form a group $\mathcal{O}(\mathcal{H})$, called the orthogonal group of X .

Denote I^+ (or I^-) to be the free Abelian group of rank 1, with basis u (or v) satisfying $u \cdot u = 1$ (or -1).

Theorem 2.1. *For $n \leq 3$, under $\mathcal{O}(I^+ + nI^-)$, each vector of positive norm is equivalent to a reduced vector $au + b_1v_1 + \cdots + b_nv_n$ with $b_1 \geq b_2 \geq \cdots \geq b_n \geq 0$, $a \geq b_1 + \cdots + b_n$.*

The theorem is obvious for $n = 1$. For $n = 2$, this is [5, Theorem 2.3]. Thus we need only to prove it for $n = 3$.

Lemma 2.2 (see [5, 1.6]). *$\mathcal{O}(I^+ + 3I^-)$ is generated by trivial automorphisms and R , where trivial automorphisms refer to permutations of v_i, v_j ; changing the signs of u and (or) v_i ; and*

$$R: \begin{cases} u \mapsto 2u + v_1 + v_2 + v_3, \\ v_1 \mapsto -u - v_2 - v_3, \\ v_2 \mapsto -u - v_1 - v_3, \\ v_3 \mapsto -u - v_1 - v_2. \end{cases}$$

here u, v_i are the basis of I^+ and I^- respectively.

Proof of Theorem 2.1 for $n = 3$. After probably changing the sign of u and v , we may assume $au + b_1v_1 + b_2v_2 + b_3v_3$ satisfy $a, b_i \geq 0$. If $a < b_1 + b_2 + b_3$, then $R(au + b_1v_1 + b_2v_2 + b_3v_3) = a'u + b'_1v_1 + b'_2v_2 + b'_3v_3$, where $a' = 2a - b_1 - b_2 - b_3 < a$.

Note that $a^2 - b_1^2 - b_2^2 - b_3^2 > 0$, so

$$\begin{aligned} 4a^2 &> 4(b_1^2 + b_2^2 + b_3^2) \\ &> (b_1^2 + b_2^2 + b_3^2) + (b_1^2 + b_2^2) + (b_2^2 + b_3^2) + (b_3^2 + b_1^2) \\ &\geq (b_1 + b_2 + b_3)^2. \end{aligned}$$

Hence

$$2a > b_1 + b_2 + b_3.$$

Hence

$$0 < a' < a.$$

Next, we induct on a . If $|a| = 1$, then $1 \geq b_1^2 + b_2^2 + b_3^2 \geq |b_1| + |b_2| + |b_3|$. For arbitrary a with $|a| < |b_1| + |b_2| + |b_3|$, we can use R on $|a|u + |b_1|v_1 + |b_2|v_2 + |b_3|v_3$ to decrease $|a|$ till we have $|a| \geq |b_1| + |b_2| + |b_3|$. Then use trivial automorphisms to get the desired vector. \square

3. Representing homology classes

Theorem 3.1. *Let $\xi, \eta_1, \eta_2, \dots, \eta_m$ be canonical generators of $H_2(\mathbb{C}P^2 \# m\overline{\mathbb{C}P}^2, \mathbb{Z})$. $u = a\xi + \sum_{i=1}^m b_i\eta_i$ has positive square, $a, b_i \in \mathbb{Z}$. We can suppose $|b_1| \geq |b_2| \geq \dots \geq |b_m|$, w.l.o.g. If $|a| \neq |b_1| + 1$ and $u^2 \geq \max\{\sum_{i=2}^m b_i^2 + 4, 5\}$, then u cannot be represented by an embedded sphere.*

Proof. Denote $n = u^2 = a^2 - \sum_{i=1}^m b_i^2$, w.l.o.g., we may assume $a > b_1 \geq b_2 \geq \dots \geq b_m \geq 0$, then $n \geq \max\{\sum_{i=2}^m b_i^2 + 4, 5\}$. Let $M = \mathbb{C}P^2 \# m\overline{\mathbb{C}P}^2 \# (n-1)\overline{\mathbb{C}P}^2$, with ζ_i , $i = 1, 2, \dots, n-1$, being the generators of $H_2(M, \mathbb{Z})$ with respect to the additional $(n-1)\overline{\mathbb{C}P}^2$. Suppose conversely that u is represented by an embedded sphere. Then $v = u + \sum_{i=1}^{n-1} \zeta_i$ can be represented by a smoothly embedded sphere S in M . The self-intersection number of $S = S \cdot S = v^2 = u^2 - (n-1) = 1$. Hence the tubular neighborhood N of S in M is the $(+1)$ -Hopf bundle over S . So ∂N is diffeomorphic to S^3 . Set $W^4 = (M^4 - \text{int } N) \cup_{\partial} D^4$. It is known [2] that W^4 is a closed, simply connected 4-manifold with negative intersection form. By Donaldson's result [1], the intersection form of W is standard. But $M = W \# \tilde{N}$, where $\tilde{N} = N \cup_{\partial} D^4 \cong \mathbb{C}P^2$. Hence the intersection form of M is the direct sum of that of W and \tilde{N} . Therefore $(H_2(W), \langle, \rangle_W) \cong (v^\perp, \langle, \rangle_M)$. So there are $2(m+n-1)$ α 's satisfying $\alpha \in H_2(W, \mathbb{Z})$, $\alpha \cdot v = 0$ and $\alpha^2 = -1$.

Write $\alpha = x\xi + \sum_{i=1}^m y_i\eta_i + \sum_{j=1}^{n-1} z_j\zeta_j$. Then the Diophantine equations

$$\begin{cases} ax = \sum_{i=1}^m b_i y_i + \sum_{j=1}^{n-1} z_j, \\ x^2 + 1 = \sum_{i=1}^m y_i^2 + \sum_{j=1}^{n-1} z_j^2 \end{cases} \quad (*)$$

shall have $2(m+n-1)$ solutions.

$$\begin{aligned} b_1^2 y_1^2 &= \left(ax - \sum_{i=2}^m b_i y_i - \sum_{j=1}^{n-1} z_j \right)^2 \\ &= b_1^2 \left(x^2 + 1 - \sum_{i=2}^m y_i^2 - \sum_{j=1}^{n-1} z_j^2 \right), \end{aligned}$$

i.e.,

$$\begin{aligned} & (a^2 - b_1^2)x^2 - 2a\left(\sum_{i=2}^m b_i y_i + \sum_{j=1}^{n-1} z_j\right)x + \left(\sum_{i=2}^m b_i y_i + \sum_{j=1}^{n-1} z_j\right)^2 \\ & + b_1^2\left(\sum_{i=2}^m y_i^2 + \sum_{j=1}^{n-1} z_j^2 - 1\right) = 0. \end{aligned}$$

We view it as a quadratic equation of x , and denote Δ to be its discriminant. Then

$$\begin{aligned} \frac{1}{4}\Delta &= a^2\left(\sum_{i=2}^m b_i y_i + \sum_{j=1}^{n-1} z_j\right)^2 - (a^2 - b^2) \\ &\quad \times \left[\left(\sum_{i=2}^m b_i y_i + \sum_{j=1}^{n-1} z_j\right)^2 + b_1^2\left(\sum_{i=2}^m y_i^2 + \sum_{j=1}^{n-1} z_j^2 - 1\right)\right] \\ &= b_1^2\left(\sum_{i=2}^m b_i y_i + \sum_{j=1}^{n-1} z_j\right)^2 - b_1^2(a^2 - b_1^2)\left(\sum_{i=2}^m y_i^2 + \sum_{j=1}^{n-1} z_j^2 - 1\right). \end{aligned}$$

Suppose $b_1 \neq 0$ and let

$$d = \frac{1}{4b_1^2}\Delta = \left(\sum_{i=2}^m b_i y_i + \sum_{j=1}^{n-1} z_j\right)^2 - (a^2 - b_1^2)\left(\sum_{i=2}^m y_i^2 + \sum_{j=1}^{n-1} z_j^2 - 1\right).$$

Denote r to be the number of z_j 's which are nonzero, s to be the number of z_j 's with $|z_j| > 1$. Then $0 \leq s \leq r \leq n-1$. After renumbering, we may suppose $z_j \neq 0$ for $1 \leq j \leq r$, and $|z_j| > 1$, for $1 \leq j \leq s$. We divide equation (*) into four cases.

Case A: $r = n - 1$. Then,

$$\begin{aligned} d &= \left(\sum_{i=2}^m b_i y_i + \sum_{j=1}^{n-1} z_j\right)^2 - (a^2 - b_1^2)\left(\sum_{i=2}^m y_i^2 + \sum_{j=1}^{n-1} z_j^2 - 1\right) \\ &\leq \left(\sum_{i=2}^m b_i |y_i| + \sum_{j=1}^s |z_j| + n - s - 1\right)^2 \\ &\quad - (a^2 - b_1^2)\left(\sum_{i=2}^m y_i^2 + \sum_{j=1}^s z_j^2 + n - s - 2\right) \\ &\quad \text{(here we use the definition of } s \text{ and the condition } r = n - 1) \\ &= \left(\sum_{i=2}^m b_i |y_i| + \sum_{j=1}^s |z_j|\right)^2 + 2(n - s - 1)\left(\sum_{i=2}^m b_i |y_i| + \sum_{j=1}^s |z_j|\right) \\ &\quad + (n - s - 1)^2 - (a^2 - b_1^2)\left(\sum_{i=2}^m y_i^2 + \sum_{j=1}^s z_j^2\right) \end{aligned}$$

$$\begin{aligned}
& - (a^2 - b_1^2)(n - s - 2) \\
& \leq \left(s + \sum_{i=2}^m b_i^2 \right) \left(\sum_{i=2}^m y_i^2 + \sum_{j=1}^s z_j^2 \right) + 2(n - s - 1) \left(\sum_{i=2}^m b_i |y_i| + \sum_{j=1}^s |z_j| \right) \\
& + (n - s - 1)^2 - \left(n + \sum_{i=2}^m b_i^2 \right) \left(\sum_{i=2}^m y_i^2 + \sum_{j=1}^s z_j^2 \right) \\
& - \left(n + \sum_{i=2}^m b_i^2 \right) (n - s - 2)
\end{aligned}$$

(here the Cauchy inequality was used for the first term)

$$\begin{aligned}
& = (s - n) \left(\sum_{i=2}^m y_i^2 + \sum_{j=1}^s z_j^2 \right) + 2(n - s - 1) \left(\sum_{i=2}^m b_i |y_i| + \sum_{j=1}^s |z_j| \right) \\
& - \left(s + \sum_{i=2}^m b_i^2 \right) n + (s + 1)^2 + (s + 2) \sum_{i=2}^m b_i^2 \\
& = \left((s - n) \sum_{j=1}^s z_j^2 + 2(n - s - 1) \sum_{j=1}^s |z_j| \right) \\
& + \left((s - n) \sum_{i=2}^m y_i^2 + 2(n - s - 1) \sum_{i=2}^m b_i |y_i| - n \sum_{i=2}^m b_i^2 \right) \\
& - sn + (s + 1)^2 + (s + 2) \sum_{i=2}^m b_i^2.
\end{aligned}$$

Note that $|z_j| \geq 2$, so $2(n - s - 1) \sum_{j=1}^s |z_j| \leq (n - s - 1) \sum_{j=1}^s z_j^2$, thus

$$\begin{aligned}
d & \leq - \sum_{j=1}^s z_j^2 - (n - s - 1) \sum_{i=2}^m (|y_i| - b_i)^2 - \sum_{i=2}^m y_i^2 - (s + 1) \sum_{i=2}^m b_i^2 - sn \\
& + (s + 1)^2 + (s + 2) \sum_{i=2}^m b_i^2 \\
& \leq -sn - (n - s - 1) \sum_{i=2}^m (|y_i| - b_i)^2 - \sum_{i=2}^m y_i^2 - 4s + (s + 1)^2 + \sum_{i=2}^m b_i^2 \\
& = -sn + (s - 1)^2 + \sum_{i=2}^m b_i^2 - (n - s - 1) \sum_{i=2}^m (|y_i| - b_i)^2 - \sum_{i=2}^m y_i^2.
\end{aligned}$$

(i) If $s \geq \sum_{i=2}^m b_i^2 \neq 0$ (we assume $s \geq 1$ if $\sum_{i=2}^m b_i^2 = 0$), then

$$\begin{aligned}
d & \leq -sn + (s - 1)^2 + \sum_{i=2}^m b_i^2 \\
& < -s(s + 1) + s^2 + \sum_{i=2}^m b_i^2 \\
& \leq 0.
\end{aligned}$$

(ii) If $1 \leq s < \sum_{i=2}^m b_i^2$, then

$$\begin{aligned} d &\leq -sn + (s-1)^2 + \sum_{i=2}^m b_i^2 \\ &< -sn + s^2 + \sum_{i=2}^m b_i^2 \\ &\leq \max \left\{ -n + 1 + \sum_{i=2}^m b_i^2, -\left(\sum_{i=2}^m b_i^2 \right) n + \left(\sum_{i=2}^m b_i^2 \right)^2 + \sum_{i=2}^m b_i^2 \right\} \\ &< 0 \end{aligned}$$

since $n \geq \sum_{i=2}^m b_i^2 + 4$.

(iii) If $s = 0$, and there exists $l \in \{2, \dots, m\}$, such that $|y_l| \neq b_l$, then

$$\begin{aligned} d &\leq -(n-1) \sum_{i=2}^m (|y_i| - b_i)^2 - \sum_{i=2}^m y_i^2 + 1 + \sum_{i=2}^m b_i^2 \\ &\leq -(n-1) + 1 + \sum_{i=2}^m b_i^2 \\ &< 0. \end{aligned}$$

(iv) If $s = 0$ and $|y_i| = b_i$ for all $i = 2, \dots, m$, we have $d \leq 1$. The equality holds iff there are no different signs among y_i, z_j for all $i = 2, \dots, m, j = 1, \dots, n-1$. Then the Diophantine equations (*) become

$$\begin{cases} x^2 + 1 = \sum_{i=2}^m b_i^2 + n - 1 + y_1^2 = a^2 - b_1^2 - 1 + y_1^2, \\ ax = \pm \left(\sum_{i=2}^m b_i^2 + n - 1 \right) + b_1 y_1 = \pm (a^2 - b_1^2 - 1) + b_1 y_1. \end{cases}$$

When $a \neq b_1 + 1$, eliminating y_1 we get

$$(a^2 - b_1^2)x^2 \pm 2a(a^2 - b_1^2 - 1)x + (a^2 - b_1^2 - 1)^2 + b_1^2(a^2 - b_1^2 - 2) = 0.$$

So

$$x = \pm a \pm \frac{a \pm b_1}{a^2 - b_1^2},$$

which is not an integer. Hence there is no solution when $a \neq b_1 + 1$.

When $a = b_1 + 1$, we have two solutions

$$\begin{cases} x = b_1, \\ y_1 = b_1 - 1, \\ y_i = b_i, \\ z_j = 1, \text{ and} \end{cases} \quad \begin{cases} x = -b_1, \\ y_1 = -b_1 + 1, \\ y_i = -b_i, \\ z_j = -1. \end{cases}$$

If there are different signs among $\{y_i, z_j | i = 2, \dots, m, j = 1, \dots, n-1\}$, it is easy to check that $d < 0$.

Case B: $2 \leq r \leq n-2$. Then

$$\begin{aligned}
 d &= \left(\sum_{i=2}^m b_i y_i + \sum_{j=1}^r z_j \right)^2 - (a^2 - b_1^2) \left(\sum_{i=2}^m y_i^2 + \sum_{j=1}^{n-1} z_j^2 - 1 \right) \\
 &\leq \left(r + \sum_{i=2}^m b_i^2 \right) \left(\sum_{i=2}^m y_i^2 + \sum_{j=1}^r z_j^2 \right) - \left(n + \sum_{i=2}^m b_i^2 \right) \left(\sum_{i=2}^m y_i^2 + \sum_{j=1}^{n-1} z_j^2 - 1 \right) \\
 &\quad (\text{here the Cauchy inequality was used and equality holds iff } b_i/y_i = z_j \text{ for all } i, j) \\
 &= (r-n) \left(\sum_{i=2}^m y_i^2 + \sum_{j=1}^r z_j^2 \right) + n + \sum_{i=2}^m b_i^2 \\
 &< r(r-n) + n + \sum_{i=2}^m b_i^2.
 \end{aligned}$$

Since $2 \leq r \leq n-2$, $r(r-n)$ reaches its maximum exactly when $r = 2$ or $n-2$. So

$$\begin{aligned}
 d &< 2(2-n) + n + \sum_{i=2}^m b_i^2 \\
 &\leq 0,
 \end{aligned}$$

since $n \geq \max\{\sum_{i=2}^m b_i^2 + 4, 5\}$.

Case C: $r = 1$. Then

$$\begin{aligned}
 d &= \left(\sum_{i=2}^m b_i y_i + z_l \right)^2 - (a^2 - b_1^2) \left(\sum_{i=2}^m y_i^2 + z_l^2 - 1 \right) \\
 &\leq \left(1 + \sum_{i=2}^m b_i^2 \right) \left(\sum_{i=2}^m y_i^2 + z_l^2 \right) - \left(n + \sum_{i=2}^m b_i^2 \right) \left(\sum_{i=2}^m y_i^2 + z_l^2 - 1 \right) \\
 &= -(n-1) \left(\sum_{i=2}^m y_i^2 + z_l^2 \right) + n + \sum_{i=2}^m b_i^2.
 \end{aligned}$$

(i) If there exists some i such that $y_i \neq 0$ or $|z_l| \neq 1$, then $d < -2(n-1) + n + \sum_{i=2}^m b_i^2 < 0$.

(ii) If $y_i = 0$ for all $i = 2, \dots, m$, and $|z_l| = 1$, then equation (*) becomes

$$\begin{cases} ax = b_1 y_1 + z_l, \\ x^2 + 1 = y_1^2 + 1, \\ z_l = \pm 1, \end{cases}$$

which has no solutions when $a \neq b_1 + 1$. When $a = b_1 + 1$, we solve it

$$\begin{cases} x = y_1 = z_l = \pm 1, \\ a = b_1 + 1, \\ y_i = 0, i = 2, \dots, m, \end{cases}$$

where l may be anyone that belongs to $\{1, \dots, n-1\}$, which has $2(n-1)$ solutions.

Case D: $r = 0$. Then

$$\begin{aligned} d &= \left(\sum_{i=2}^m b_i y_i \right)^2 - (a^2 - b_1^2) \left(\sum_{i=2}^m y_i^2 - 1 \right) \\ &\leq \left(\sum_{i=2}^m b_i^2 \right) \left(\sum_{i=2}^m y_i^2 \right) - \left(n + \sum_{i=2}^m b_i^2 \right) \left(\sum_{i=2}^m y_i^2 - 1 \right) \\ &= -n \sum_{i=2}^m y_i^2 + n + \sum_{i=2}^m b_i^2. \end{aligned}$$

(i) If $\sum_{i=2}^m y_i^2 \geq 2$, then

$$d \leq -2n + n + \sum_{i=2}^m b_i^2 < 0.$$

(ii) If $\sum_{i=2}^m y_i^2 = 1$, i.e., there is some $l \in \{2, \dots, m\}$ such that $|y_l| = 1$ and $y_i = 0$ for all $i \neq l$. Then equation (*) becomes

$$\begin{cases} x^2 + 1 = y_1^2 + 1, \\ ax = b_1 y_1 + b_l y_l, \\ y_l = \pm 1. \end{cases}$$

That is

$$\begin{cases} x = y_1, z_j = 0, j = 1, \dots, n-1, \\ y_l = \pm 1, l \in \{2, \dots, m\}, \\ y_i = 1, i \in \{2, \dots, m\} \text{ and } i \neq l, \\ (a - b_1)x = b_l y_l, \end{cases}$$

which has at most $2(m-1)$ solutions.

(iii) If $\sum_{i=2}^m y_i^2 = 0$, i.e., $y_i = 0$ for $i = 2, \dots, m$, then (*) becomes

$$\begin{cases} x^2 + 1 = y_1^2, \\ ax = b_1 y_1, \end{cases}$$

which has no solutions.

Summarizing the above deductions, we see that if $a \neq b_1 + 1$ and $n \geq \sum_{i=2}^m b_i^2 + 4$ or 5, then the Diophantine equations (*) have fewer than $2(n+m-1)$ solutions.

The above discussions are based on $b_1 \neq 0$. If $b_1 = 0$, this means $b_i = 0$ for all $i = 1, \dots, m$, then (*) becomes

$$\begin{cases} ax = \sum_{j=1}^{n-1} z_j, \\ x^2 + 1 = \sum_{i=2}^m y_i^2 + \sum_{j=1}^{n-1} z_j^2, \quad n = a^2. \end{cases}$$

So, we use the Cauchy inequality

$$a^2 x^2 = \left(\sum_{j=1}^r z_j \right)^2 \leq r \sum_{j=1}^r z_j^2 \leq r(x^2 + 1).$$

Hence

$$\begin{aligned} x^2 &\leq r, \\ \sum_{i=1}^m y_i^2 + \sum_{j=1}^r z_j^2 &\leq r + 1. \end{aligned}$$

This implies $|z_j| = 1$ for $j = 1, \dots, r$ and $\sum_{i=1}^m y_i^2 \leq 1$.

If $\sum_{i=1}^m y_i^2 = 1$, then

$$\begin{cases} x^2 + 1 = 1 + \sum_{j=1}^r z_j^2 = 1 + r, \\ ax = r - 2r', \end{cases}$$

where r' is the number of z_j 's with $z_j = -1$.

$$a^2 x^2 = (r - 2r')^2,$$

so

$$a^2 r = (r - 2r')^2 \leq r^2 \leq (a^2 - 1)r.$$

Therefore

$$r = 0.$$

In this case there are $2m$ solutions:

$$\begin{cases} x = z_j = 0, & j = 1, \dots, n-1, \\ y_l = \pm 1, & l \in \{1, \dots, m\}, \\ y_i = 0, & \text{for all } i \neq l. \end{cases}$$

If $\sum_{i=1}^m y_i^2 = 0$, then

$$\begin{aligned} ax &= r - 2r', \\ x^2 + 1 &= r, \\ a^2 x^2 &= (r - 2r')^2 = a^2(r - 1). \end{aligned}$$

Note $r \leq n - 1 = a^2 - 1$, so we have $r' = 0$. Therefore,

$$x^2 + 1 = ax = r,$$

$$x^2 - ax + 1 = 0,$$

which has integer solutions iff $a = \pm 2$.

Thus the theorem is proved. \square

From the proof, we can also see

Theorem 3.2. $u = a\xi \in H_2(\mathbb{C}P^2 \# m\overline{\mathbb{C}P}^2, \mathbb{Z})$ is represented by an embedded sphere iff $|a| \leq 2$, where ξ is the canonical generator of $H_2(\mathbb{C}P^2 \# m\overline{\mathbb{C}P}^2, \mathbb{Z})$ with respect to $\mathbb{C}P^2$.

In Section 2, we proved for $m \leq 3$, each element in $H_2(\mathbb{C}P^2 \# m\overline{\mathbb{C}P}^2, \mathbb{Z})$ with positive square is equivalent under some automorphisms to a reduced vector $au + b_1v_1 + \cdots + b_mv_m$ with $b_1 \geq b_2 \geq \cdots \geq b_m \geq 0$ and $a \geq b_1 + b_2 + \cdots + b_m$. Theorem 2 of Wall [6] says every such automorphism is induced by a diffeomorphism of M . Thus for $n = 2$ or 3 , we need only to consider such reduced vectors and this yields:

Theorem 3.3. Let $u = (a, b_1, b_2) \in H_2(\mathbb{C}P^2 \# 2\overline{\mathbb{C}P}^2, \mathbb{Z})$ satisfy $|b_1| \geq |b_2|$ and $|a| \geq |b_1| + |b_2|$. Then u is represented by an embedded sphere iff $|a| - |b_1| \leq 1$ or $u = (\pm 2, 0, 0)$.

Theorem 3.4. Let $u = (a, b_1, b_2, b_3) \in H_2(\mathbb{C}P^2 \# 3\overline{\mathbb{C}P}^2, \mathbb{Z})$ satisfy $|b_1| \geq |b_2| \geq |b_3|$ and $|a| \geq |b_1| + |b_2| + |b_3|$. Then u is represented by an embedded sphere iff $b_3 = 0$ and $|a| - |b_1| \leq 1$ or $u = (\pm 2, 0, 0, 0)$.

The if part is known (see Wall [6] or Lawson [3]). The other part of Theorems 3.3 and 3.4 is direct from Theorem 3.2.

As a consequence we can deduce Lawson's theorem on representing homology classes in $\mathbb{C}P^2 \# \overline{\mathbb{C}P}^2$ and that of Kuga on $S^2 \times S^2$ from our results.

Corollary 3.5 (Lawson and Luo). $u = ax + by \in H_2(\mathbb{C}P^2 \# \overline{\mathbb{C}P}^2, \mathbb{Z})$ is represented by an embedded sphere iff $||a| - |b|| \leq 1$ or $(a, b) = (0, \pm 2)$ or $(\pm 2, 0)$.

Proof. The if part is known (see Wall [6]). Suppose $||a| - |b|| \geq 2$, w.l.o.g. we may suppose $a - 2 \geq b \geq 0$. If (a, b) is represented by an embedded sphere in $\mathbb{C}P^2 \# \overline{\mathbb{C}P}^2$, then $(a, b, 1)$ is represented in $\mathbb{C}P^2 \# \overline{\mathbb{C}P}^2 \# \overline{\mathbb{C}P}^2$, which contradicts to Theorem 3.3, since $a \geq b + 2$. \square

Corollary 3.6 (Kuga). $p\xi + q\eta \in H_2(S^2 \times S^2, \mathbb{Z})$ is represented by an embedded sphere iff $|p| \leq 1$ or $|q| \leq 1$.

Proof. The if part is trivial. Suppose w.l.o.g. $p \geq 2$, $q \geq 2$ and $p\xi + q\eta$ is represented by an embedded sphere. Then $p\xi + q\eta + \zeta$ is represented in $S^2 \times S^2 \# \overline{\mathbb{C}P}^2$, where ζ is the canonical generator of $H_2(\overline{\mathbb{C}P}^2)$. Note $S^2 \times S^2 \# \overline{\mathbb{C}P}^2$ is diffeomorphic to $\mathbb{C}P^2 \# \overline{\mathbb{C}P}^2 \# \overline{\mathbb{C}P}^2$. Let $\xi = u - v_1$, $\eta = u - v_2$, $\zeta = v_1 + v_2 - u$. Then $u^2 = -v_1^2 = -v_2^2 = 1$, $uv_1 = uv_2 = v_1v_2 = 0$. Hence u, v_1, v_2 can be viewed as obtained from the canonical generators of $H_2(\mathbb{C}P^2 \# \overline{\mathbb{C}P}^2 \# \overline{\mathbb{C}P}^2, \mathbb{Z})$ through an automorphism. $p\xi + q\eta + \zeta = (p + q - 1)u - (p - 1)v_1 - (q - 1)v_2$. By Theorem 3.3, it is not represented by a smoothly embedded sphere. A contradiction. \square

Note in $\mathbb{C}P^2 \# \overline{\mathbb{C}P}^2$, if $\|a\| - \|b\| = 1$, then $a\xi + b\eta \in H_2(\mathbb{C}P^2 \# \overline{\mathbb{C}P}^2, \mathbb{Z})$ is represented by an embedded sphere, where ξ, η are the standard generators of $H_2(\mathbb{C}P^2 \# \overline{\mathbb{C}P}^2, \mathbb{Z})$. The next theorem shows a similar result.

Theorem 3.7. Let $u = a\xi + \sum_{i=1}^m b_i \eta_i \in H_2(\mathbb{C}P^2 \# m\overline{\mathbb{C}P}^2, \mathbb{Z})$, where ξ, η_i are the canonical generators of $H_2(\mathbb{C}P^2 \# m\overline{\mathbb{C}P}^2, \mathbb{Z})$. If $\|a\| - \|b_i\| = 1$ for some $i \in \{1, \dots, m\}$, then u can be represented by an embedded sphere.

Proof. We induct on m . For $m = 2$, w.l.o.g. we may assume $\|a\| - \|b_1\| = 1$. Change the sign of generators if necessary, we can make a, b_1, b_2 positive. Then $a = b_1 + k$, $k = 1$ or -1 . It is easy to check that

$$T_1: \begin{cases} \xi \rightarrow 3\xi + 2\eta_1 + 2\eta_2, \\ \eta_1 \rightarrow -2\xi - \eta_1 - 2\eta_2, \\ \eta_2 \rightarrow -2\xi - 2\eta_1 - \eta_2 \end{cases}$$

is an automorphism of $H_2(\mathbb{C}P^2 \# 2\overline{\mathbb{C}P}^2, \mathbb{Z})$.

$$\begin{aligned} (b_1 + k, b_1, b_2) &\xrightarrow{T_1} (b_1 + 3k - 2b_2, b_1 + 2k - 2b_2, 2k - b_2) \\ &\xrightarrow{T_2} (b_1 + 3k - 2b_2, b_1 + 2k - 2b_2, b_2 - 2k) \end{aligned}$$

where T_2 is the automorphism changing of the sign of η_2 . Set $T = T_2 \cdot T_1$, then

$$(b_1 + k, b_1, b_2) \xrightarrow{T} (b_1^{(1)} + k, b_1^{(1)}, b_2^{(1)})$$

where $b_1^{(1)} = b_1 + 2k - 2b_2$, $b_2^{(1)} = b_2 - 2k$.

Denote $b_i^{(n)} = T^n b_i$, $i = 1, 2$. So we have,

$$\begin{cases} b_2^{(n+1)} = b_2^{(n)} - 2k, \\ b_1^{(n+1)} = b_1^{(n)} + 2k - 2b_2^{(n)}. \end{cases}$$

Hence

$$\begin{cases} b_2^{(n)} = b_2 - 2kn, \\ b_1^{(n)} = b_1 - 2b_2n + 2kn^2. \end{cases}$$

We can choose n , such that $0 \leq b_2 - 2n \leq 1$.

If $k = 1$, then $(b_1 + 1, b_1, b_2) \xrightarrow{T^n} (b_1^{(n)} + 1, b_1^{(n)}, b_2^{(n)})$ which is obviously represented by an embedded sphere, since $b_2^{(n)} = 0$ or 1 . By Theorem 2 of Wall [6], $(b_1 + 1, b_1, b_2) = (a, b_1, b_2)$ is represented by a smoothly embedded sphere. If $k = -1$, set $T' = T_1 \circ T_2$, then $(b_1 - 1, b_1, b_2) \xrightarrow{T'^n} (b_1^{(n)} - 1, b_1^{(n)}, b_2^{(n)})$, where $b_i^{(n)} = T'^n b_i$ and $b_2^{(n)} = b_2 + 2kn = b_2 - 2n = 0$ or 1 . Hence $(b_1 - 1, b_1, b_2) = (a, b_1, b_2)$ is represented by an S^2 . Thus the theorem holds for $m = 2$. Suppose the theorem holds for $m (\geq 2)$. For $m + 1$, we suppose $||a| - |b_1|| = 1$ and a, b_1, b_2 positive. From the deduction above, we see that

$$(a, b_1, b_2, b_3, \dots, b_{m+1}) \xrightarrow{R^n \oplus \text{Id}} (a^{(n)}, b_1^{(n)}, b_2^{(n)}, b_3, \dots, b_{m+1})$$

where $R = T$ or T' depending on $a - b_1 = 1$ or -1 respectively, n is a positive integer satisfying $0 \leq b_2 - 2n \leq 1$. Since $|a^{(n)} - b_1^{(n)}| = 1$, by the inductive hypothesis, $(a^{(n)}, b_1^{(n)}, b_3, \dots, b_{m+1})$ is representable in $\mathbb{C}P^2 \# m \overline{\mathbb{C}P}^2$. Note $b_2^{(n)} = 0$ or 1 , therefore $(a^{(n)}, b_1^{(n)}, b_2^{(n)}, b_3, \dots, b_{m+1})$ is representable in $\mathbb{C}P^2 \# (m+1) \overline{\mathbb{C}P}^2$ by an embedded S^2 . By Theorem 2 of Wall [6] the automorphism $R^n : H_2(\mathbb{C}P^2 \# 2 \overline{\mathbb{C}P}^2, \mathbb{Z}) \rightarrow H_2(\mathbb{C}P^2 \# 2 \overline{\mathbb{C}P}^2, \mathbb{Z})$ is induced by a diffeomorphism $f : \mathbb{C}P^2 \# 2 \overline{\mathbb{C}P}^2 \rightarrow \mathbb{C}P^2 \# 2 \overline{\mathbb{C}P}^2$. Then, $R^n \oplus \text{Id}$ is induced by the diffeomorphism $f \# \text{Id} : \mathbb{C}P^2 \# 2 \overline{\mathbb{C}P}^2 \# (m-1) \overline{\mathbb{C}P}^2 \rightarrow \mathbb{C}P^2 \# 2 \overline{\mathbb{C}P}^2 \# (m-1) \overline{\mathbb{C}P}^2$, where Id is the identity map of $(m-1) \overline{\mathbb{C}P}^2$ onto itself. This proves $(a, b_1, b_2, b_3, \dots, b_{m+1})$ is represented by an embedded sphere, and hence the theorem. \square

Remark 3.8. Using a similar technique we can show that a vector $(a + 2, a, b) \in H_2(\mathbb{C}P^2 \# 2 \overline{\mathbb{C}P}^2, \mathbb{Z})$ with positive square is represented by an embedded S^2 iff $a = n^2$, $b = 2n$ or $a = 4n^2 + 2n$, $b = 4n + 1$.

Remark 3.9. After submitting this paper the author became aware that a similar result was also obtained by Mr. K. Kikuchi and D.Y. Gan.

References

- [1] S.K. Donaldson, Self-dual connections and the topology of smooth 4-manifolds, *Bull. Amer. Math. Soc. (N.S.)* 8 (1983) 81–83.
- [2] K. Kuga, Representing homology classes of $S^2 \times S^2$, *Topology* 23 (1984) 133–137.
- [3] T. Lawson, Representing homology classes of almost definite 4-manifolds, *Michigan Math. J.* 34 (1987) 85–91.
- [4] F. Luo, Representing homology classes for $CP^2 \# \overline{CP}^2$, *Pacific J. Math.* 133 (1988) 137–140.
- [5] C.T.C. Wall, On the orthogonal groups of unimodular quadratic forms II, *Crelle J.* 213 (1963) 122–126.
- [6] C.T.C. Wall, Diffeomorphisms of 4-manifolds, *J. London Math. Soc.* 39 (1964) 131–140.